

IMPACT OF EXERCISE ON SKELETAL MUSCLE IN HEALTH AND METABOLIC DISEASES

Prof. Dr. Wook Song
Department of Kinesiology,
Institute of Sport Science,
Institute on Aging,
Seoul National University,
Seoul, South Korea.
E-mail: songw3@snu.ac.kr

Aerobic training (endurance training) has two major goals. First of all, aerobic training induces adaptation in cardiovascular function increasing oxygen delivery to active skeletal muscle. Secondly, aerobic training induces significant adaptations in muscle metabolism. This beneficial adaptation to training in skeletal muscle is accompanied by structural and biochemical change including muscle fiber type, mitochondrial number, oxidative enzymes, and energy stores and substrate availability. However, adaptations can be quickly lost with detraining. One of the impacts of exercise on skeletal muscle is an increase of resting (basal) metabolic rate. Resting metabolic rate is highly correlated with free-fat mass and significantly decreased with aging. Many studies have indicated that regular resistance exercise offsets the decrease in resting metabolism that usually accompanies aging. This is a reason why resistance exercise is important for obesity intervention because resting metabolic rate is accounting 60-75% of total daily energy expenditure.

Aging in skeletal muscle is characterized by loss of skeletal muscle mass and function (i.e., sarcopenia) along with increased susceptibility to oxidative stress, injury, and inflammation. Sarcopenia adversely affects the quality of life and independence of the elderly. Recently, sarcopenic obesity become the interesting area of research since obesity (i.e., visceral fat) is epidemic with age, and many studies were reported regarding definition, etiology and consequences of sarcopenic obesity. In collaboration with endocrinology research group, we firstly reported the prevalence of sarcopenia and sarcopenic obesity in Korean population. Strong evidences demonstrated that exercise would be the only and the best physiological intervention for delay and prevention of sarcopenia. In the presentation, the major interesting findings will be introduced and discussed from the Korean Sarcopenic Obesity Study (KSOS) primarily designed to investigate criteria and mechanisms of sarcopenic obesity related to metabolic syndrome in Korean population.

The prevalence of type 2 diabetes mellitus (DM) is dramatically increasing in recent 20 years. Type 2 DM is becoming one of the leading causes of death in Korea and is considered to result from a gradual development of insulin resistance and deterioration of the ability to transport glucose from circulating blood cross primarily muscle, liver, or other tissue cells. Growing scientific evidences suggest that the major driver in the pathogenesis of type 2 DM is inactivity of the skeletal muscle. However, it is well known that skeletal muscle increases its insulin sensitivity for glucose uptake from the blood during exercise (contraction) and even during post-exercise just long enough to replenish the stores of intramuscular glycogen consumed by exercise. In collaboration with veterinary medicine research group, we've been able to use type 2 DM animal model investigating effects of aerobic and resistance training on this metabolic disease. In this presentation, previous and current studies in our laboratory with respect to exercise and type 2 DM using animal models including Otsuka Long-Evans Tokushima Fatty (OLETF) rats and Zucker Diabetic Fatty (ZDF) rats will be introduced and discussed briefly.

References:

1. Kim TN, Yang SJ, Yoo HJ, Lim KI, Kang HJ, **Song W**, Seo JA, Kim SG, Kim NH, Baik SH, Choi DS, and Choi KM. Prevalence of sarcopenia and sarcopenic obesity in Korean adults: the Korean sarcopenic obesity study. *Int J Obes (Lond)*. 2009 Aug;33(8):885-92
2. Chang H, Park JY, Suk MH, Lee HJ, Kang HJ, Choi KM, and **Song W**. Comparison of lactate threshold, glucose, and insulin levels between OLETF and LETO rats after all-out exercise. *Journal of Sports Science and Medicine*. Sep;8(3):381-387, 2009
3. D-I. Seo, T-W. Jun, K-S. Park, H-K. Chang, W-Y. So, and **W. Song**. 12 weeks of combined exercise training is better than aerobic exercise training for increasing growth hormone in middle-aged women. *International Journal of Sports Nutrition and Exercise Metabolism*.2010 Feb;20(1):21-6
4. Kim TN, Park MS, Yang SJ, Yoo HJ, Kang HJ, **Song W**, Seo JA, Kim SG, Kim NH, Baik SH, Choi DS, Choi KM. Prevalence and determinant factors of sarcopenia in patients with type 2 diabetes: the Korean Sarcopenic Obesity Study (KSOS). *Diabetes Care*. 2010 Jul;33(7):1497-9
5. Lim KI, Yang SJ, Kim TN, Yoo HJ, Kang HJ, **Song W**, Baik SH, Choi DS, Choi KM. The association between the ratio of visceral fat to thigh muscle area and

- metabolic syndrome: the Korean Sarcopenic Obesity Study (KSOS). *Clin Endocrinology (Oxf)*. 2010 Nov;73(5):588-94
6. K-J. Lee, Y-A. Shin, K-Y. Lee, T-W. Jun, and **W. Song**. Aerobic exercise training-induced decrease in plasma visfatin and insulin resistance in obese female adolescents. *International Journal of Sport Nutrition and Exercise Metabolism*. 2010 Aug;20(4):275-81
 7. Lee H, Chang H, Park JY, Kim SY, Choi KM, **Song W**. Exercise training improves basal blood glucose metabolism with no changes of cytosolic IKK, JNK activation in skeletal muscle of OLETF rat. *Exp Physiol*. 2011 Jul; 96(7):689-98
 8. Kim TN, Park MS, Lim KI, Yang SJ, Yoo HJ, Kang HJ, **Song W**, Seo JA, Kim SG, Kim NH, Baik SH, Choi DS, Choi KM. Skeletal muscle mass to visceral fat area ratio is associated with metabolic syndrome and arterial stiffness: The Korean Sarcopenic Obesity Study (KSOS). *Diabetes Res Clin Pract*. 2011 Aug;93(2):285-91. Epub 2011 Jul 14.
 9. Song M, Park YH, **Song W**, Cho BL, Lim JY, Kim S, Choi S. Combined exercise training and self-management education for community-dwelling older adults with diabetes in Korea. *J Gerontol Nurs*. 2012 Oct; 38(10):38-48
 10. Kim TN, Park MS, Yang SJ, Yoo HJ, Kang HJ, **Song W**, Seo JA, Kim SG, Kim NH, Baik SH, Choi DS, Choi KM. Body size phenotypes and low muscle mass: the Korean sarcopenic obesity study (KSOS). *J Clin Endocrinol Metab*. 2013 Feb;98(2):811-7. Epub 2013 Jan 4.
 11. Kim HJ, Park JY, Oh SL, Kim YA, So BH, **Song W**. Effect of treadmill exercise on interleukin-15 expression and glucose tolerance in Zucker diabetic fatty rats. *Diabetes Metab J*. 2013 Oct;37(5):358-64
 12. Byunghun So, Hee-jae Kim, Jinsoo Kim, **Wook Song**. Exercise-induced myokines in health and metabolic disease. *Integrative medicine research*. 2014 Oct [online 5]
 13. Kim JY, Choi MJ, So B, Kim HJ, Seong JK, **Song W**. The Preventive Effects of 8 Weeks of Resistance Training on Glucose Tolerance and Muscle Fiber Type Composition in Zucker Rats. *Diabetes Metab J*. 2015 Oct;39(5):424-33.
 14. Kim HJ, So B, Choi M, Kang D, **Song W**. Resistance exercise training increases the expression of irisin concomitant with improvement of muscle function in aging mice and humans. *Exp Gerontol*. 2015 Oct;70:11-7.
 15. H Kim, HJ Lee, BH So, JS Son, Dh Yoon, **W Song**. Effect of aerobic training and resistance training on circulating irisin level and their association with change of

body composition in overweight/obese adults: a pilot study. *Physiol. Res.* 65: 271-279, 2016.

16. Su Han Cho, Jang Hoe Kim, **Wook Song**. In vivo rodent models of skeletal muscle adaptation to decreased use. *Endocrinol Metab (Seoul)*.2016 Mar;31(1):31-7
17. Son JS, Kim HJ, Son Y, Lee H, Chae SA, Seong JK, **Song W**. Effects of exercise-induced apelin levels on skeletal muscle and their capillarization in type 2 diabetic rats. *Muscle Nerve*. 2017 Feb 6
18. DH Yoon, DH Kang, HJ Kim, JS Kim, HS Song and **W Song**. Effect of elastic band-based high-speed power training on cognitive function, physical performance and muscle strength in older women with mild cognitive impairment. *Geriatr Gerontol Int* 2017 May;17(5):765-772.